

**Central Bureau for Astronomical Telegrams  
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)  
CBAT@CFA.HARVARD.EDU (science)  
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304  
Phone 617-495-7440/7244/7444 (for emergency use only)

*COMET P/2004 V5 (LINEAR-HILL)*

Z. Sekanina, Jet Propulsion Laboratory, writes that his comet fragmentation model (cf. *IAUC* 8434) shows that the two nuclei of P/2004 V5 (*MPECs* 2004–V79 and 2004–W07) broke apart around  $2001.9 \pm 0.3$  year, at a heliocentric distance of  $\sim 6.3$  AU and 2.5 years before perihelion. The separation velocity of the companion (fragment B) relative to the primary (A) pointed below the orbital plane and was at least 2.6 m/s. The motion of B has since been subjected to a differential deceleration of  $40 \pm 6$  units of  $10^{-5}$  the solar attraction. Predicted separations and position angles of B relative to A are as follows (equinox 2000.0): 2004 Nov. 21.0 TT,  $131''$ ,  $287^\circ$ ; Dec. 1.0,  $137''$ ,  $288^\circ$ ; 11.0,  $144''$ ,  $289^\circ$ ; 21.0,  $152''$ ,  $290^\circ$ ; 31.0,  $159''$ ,  $290^\circ$ ; 2005 Jan. 10.0,  $167''$ ,  $291^\circ$ ; 20.0,  $175''$ ,  $291^\circ$ ; 30.0,  $181''$ ,  $291^\circ$ .

*SUPERNOVA 2004fz IN NGC 783*

A. V. Filippenko and R. J. Foley, University of California, Berkeley, report that inspection of CCD spectra (range 320–730 nm), obtained by S. Mochmacki and J. Thomson (University of Toronto) on Nov. 15 UT with the 1.88-m reflector at David Dunlap Observatory, suggests that SN 2004fz (*IAUC* 8437) is of type Ia, roughly 1 week prior to maximum brightness. This classification was confirmed by inspection of higher-quality CCD spectra (range 310–710 nm) obtained by D. Kirkman and D. Tytler (University of California, San Diego) on Nov. 16 with the Shane 3-m reflector at Lick Observatory. Its spectral-feature age (Riess *et al.* 1997, *A.J.* **114**, 722), derived from the Lick data, is  $5 \pm 2$  days before maximum brightness. The expansion velocity of the ejecta, as measured from the minimum in the Si II 635.5-nm absorption trough, is  $\sim 11000$  km/s. No interstellar Na I D absorption lines are present, to an equivalent width limit of  $\sim 0.02$  nm.

*COMET C/2004 Q2 (MACHHOLZ)*

Visual total-magnitude estimates: Nov. 6.79 UT, 7.2 (A. Pearce, Noble Falls, W. Australia,  $8 \times 40$  binoculars); 13.10, 6.9 (J. J. Gonzalez, Leon, Spain,  $7 \times 50$  binoculars); 14.45, 6.2 (D. A. J. Seargent, Cowra, N.S.W., naked eye); 16.60, 6.5 (Y. Nagai, Nagano, Japan,  $7 \times 35$  binoculars).

*COMET C/2003 K4 (LINEAR)*

Visual total-magnitude estimates by Pearce ( $20 \times 80$  binoculars): Aug. 8.47 UT, 6.9; Oct. 22.84, 7.4; Nov. 6.80, 7.3; 13.80, 7.2; 17.80, 7.2.