

**Central Bureau for Astronomical Telegrams  
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)  
CBAT@CFA.HARVARD.EDU (science)  
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304  
Phone 617-495-7440/7244/7444 (for emergency use only)

*SUPERNOVA 2005K IN NGC 2923*

Further to *IAUC* 8467, O. Trondal and M. Schwartz report the discovery of an apparent supernova (mag  $\sim 17.8$ ) on unfiltered Tenagra II 0.81-m telescope images taken on Jan. 15.41 and 17.20 UT. SN 2005K is located at  $\alpha = 9^{\text{h}}36^{\text{m}}04^{\text{s}}.31$ ,  $\delta = +16^{\circ}45'46''.8$  (equinox 2000.0), which is  $6''.7$  east and  $9''.5$  north of the nucleus of NGC 2923. Nothing is visible at this position on an image taken on 2004 Sept. 12.55 (limiting mag  $\sim 19.0$ ).

*SUPERNOVAE 2004gx, 2004gz, 2005C, AND 2005G*

M. Ganeshalingam, F. J. D. Serduke, and A. V. Filippenko, University of California, Berkeley, report that inspection of CCD spectra (range 330–1060 nm), obtained on Jan. 16 UT with the Shane 3-m reflector at Lick Observatory, reveals that SN 2004gz (*IAUC* 8460) is of type Ia, a few weeks past maximum brightness. SN 2005C (*IAUC* 8463) is probably of type Ib, several weeks past maximum; a set of He I absorption lines appears to be present. SN 2005G (*IAUC* 8465) is probably of type Ia near maximum, but the spectrum is somewhat peculiar (e.g., unusually narrow Si II 635.5-nm absorption; the two S II absorption lines near 550 nm are blended together); although the optical continuum is very blue, there is a sharp decline in flux density shortward of the Ca II H + K trough. A very noisy CCD spectrum, obtained on Jan. 17, suggests that SN 2004gx (*IAUC* 8459) is probably of type II. A broad emission feature at the wavelength of H $\alpha$  is visible in the redshifted spectrum.

H. Navasardyan, S. Benetti, N. Elias-Rosa, and A. Harutunyan, Osservatorio Astronomico di Padova; and A. Pastorello, Max-Planck-Institut für Astrophysik, Garching, on behalf of the European RTN collaboration (cf. *IAUC* 7987), report that a spectrum of SN 2005G (cf. *IAUC* 8465), obtained on Jan. 18.16 UT with the Asiago 1.8-m telescope (+ AFOSC; range 355–780 nm, resolution 2.4 nm), shows it to be a type-Ia supernova  $\sim 10$  days past maximum. The spectrum closely resembles that of SN 1994D (Patat *et al.* 1996, *MNRAS* **278**, 111) at comparable phase. The expansion velocity deduced from the Si II 635.5-nm minimum is  $\sim 9400$  km/s (adopting the NED recession velocity of 6938 km/s for the parent galaxy).

*COMET 163P/2004 V4 (NEAT)*

Following the identification of observations of comet P/2004 V4 (cf. *IAUC* 8429, 8438) in 1990–1991 and 1997 (cf. *MPC* 53257, 53303, 53307), the comet has been numbered 163P.