

**Central Bureau for Astronomical Telegrams  
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)  
CBAT@CFA.HARVARD.EDU (science)  
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304  
Phone 617-495-7440/7244/7444 (for emergency use only)

*SUPERNOVAE 2005N, 2005O, AND 2005P*

Further to *IAUC* 8470, J. Burket and W. Li report the LOSS discovery of an apparent supernova on unfiltered KAIT images taken on Jan. 21.55 (at mag 18.1) and 22.54 UT (mag 18.0). SN 2005P is located at  $\alpha = 14^{\text{h}}06^{\text{m}}34^{\text{s}}.01$ ,  $\delta = -5^{\circ}27'42''.6$  (equinox 2000.0), which is  $12''.6$  west and  $31''.7$  south of the nucleus of NGC 5468. A KAIT image taken on 2004 July 8.21 showed nothing at this position (limiting mag 19.5).

B. Schmidt and M. Salvo, Australian National University, report that a spectrogram (range 420–1000 nm) of SN 2005O (cf. *IAUC* 8471), obtained by P. Wood on Jan. 21.70 UT with the ANU 2.3-m telescope (+ DBS), reveals it to be a type-Ib supernova near maximum light. The relatively red spectrum is dominated by He I lines and the Ca II infrared triplet.

S. Taubenberger and A. Pastorello, Max-Planck-Institut für Astrophysik, Garching, on behalf of the European RTN collaboration, report that inspection of a low-*S/N* spectrogram of SN 2005N (cf. *IAUC* 8470), taken on Jan. 22.21 UT by M. Alises with the Calar Alto 2.2-m telescope (+ CAFOS; range 330–880 nm), shows it to be a type-Ib/c supernova in the nebular phase,  $\sim 4$  months after the explosion. The spectrum is characterized by strong emission lines of [O I]; features due to [Ca II], Mg I], [Fe II], Na I, O I, and Ca II are also visible.

*SUPERNOVA 2001em IN UGC 11794*

C. J. Stockdale and B. Kaster, Marquette University; L. O. Sjouwerman and M. P. Rupen, National Radio Astronomy Observatory; I. Martí-Vidal and J. M. Marcaide, University of Valencia; S. D. Van Dyk, Spitzer Science Center, California Institute of Technology; K. W. Weiler, Naval Research Laboratory; B. Paczynski, Princeton University; and N. Panagia, European Space Agency and Space Telescope Science Institute, report a test of the proposal by Paczynski (2001, *Acta Astron.* **51**, 1) and by Granot and Loeb (2003, *Ap.J.* **593**, L81) that type-Ib/c supernovae may produce late-time radio jet emission. A Very Long Baseline Array measurement on 2004 July 1 of the oldest known radio-emitting type-Ib/c supernova, SN 2001em, at age  $\sim 1020$  days after explosion, shows no extended radio emission to a  $3\sigma$  limit of 0.3 mJy/beam. The detected radio emission at 8.4 GHz of  $1.8 \pm 0.2$  mJy from SN 2001em appears unresolved at a resolution of  $0''.0019 \times 0''.0008$ , corresponding to a radius of  $< 0.4 \pm 0.2$  pc at a distance of 90 Mpc or an average expansion speed for any detectable radio emitting material of  $< 150000$  km/s ( $< 0.5c$ ).