

**Central Bureau for Astronomical Telegrams  
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)  
CBAT@CFA.HARVARD.EDU (science)  
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304  
Phone 617-495-7440/7244/7444 (for emergency use only)

*2004 MN<sub>4</sub>*

On 2004 Dec. 20, K. E. Smalley (cf. *MPEC* 2004-Y25) identified an object placed on the ‘NEO Confirmation Page’ on Dec. 18 (following its discovery by G. J. Garradd via the Siding Spring Survey) with 2004 MN<sub>4</sub> (which had been recorded at Kitt Peak on June 19 and 20; cf. *MPS* 109613). Although the recognition of further predisccovery observations by Spacewatch on Mar. 15 (*MPEC* 2004-Y70) precluded the possible 2029 Apr. 13 earth impact discussed extensively in the WWW during 2004 Dec. 23–27 [notably on the Jet Propulsion Laboratory (JPL) and Pisa NEODys “risk pages”], it was clear that the object would then make an unusually close approach. L. A. M. Benner, JPL; M. C. Nolan, National Astronomy and Ionosphere Center, Arecibo Observatory; J. D. Giorgini, S. R. Chesley, and S. J. Ostro, JPL; and D. J. Scheeres, University of Michigan, report: “Arecibo delay-Doppler radar astrometry obtained on 2005 Jan. 27, 29, and 30 significantly refines the 2004 MN<sub>4</sub> orbit. On Jan. 29.0 UT, the range was 294 km closer to the earth than the pre-radar orbit solution predicted. This correction results in a 2029 approach to the geocenter of only  $0.000245 \pm 0.000060$  AU ( $36700 \pm 9000$  km or  $5.7 \pm 1.4$  earth radii,  $3\sigma$  uncertainties), which is just below geosynchronous orbit and 28000 km closer than predicted by the pre-radar ephemeris. During its close approach, it is likely that tidal torques will significantly alter the object’s spin state.”

*ARP 299*

Further to *IAUC* 8473, S. Mattila, R. Greimel, C. Gerardy, and W. P. S. Meikle, together with D. L. Clements and K. Nandra (Imperial College, London), report the discovery on a  $K_s$ -band image obtained on Jan. 30.3 UT of a strong outburst in the B1 nucleus (Wynn-Williams *et al.* 1991, *Ap.J.* **377**, 426) of the galaxy Arp 299. The nucleus was not in outburst in their previous such image taken on 2004 June 6.0, and the difference in brightness yields a magnitude for the new source of  $K_s = +12.6 \pm 0.1$ ; at a distance of 45 Mpc, this translates to  $M(K_s) \sim -20.7$ , which is about a magnitude brighter than would be expected for the brightest supernovae (cf. Mattila and Meikle 2001, *MNRAS* **324**, 325). The new source is located at  $\alpha = 11^{\text{h}}28^{\text{m}}31^{\text{s}}02$ ,  $\delta = +58^{\circ}33'40''.7$  (equinox 2000.0), which is  $< 0''.1$  from the  $K_s$ -band nucleus B1 and  $< 0''.5$  from the location of the x-ray-determined active galactic nucleus (AGN) in Arp 299 (Ballo *et al.* 2004, *Ap.J.* **600**, 634). The source is therefore most likely associated with an AGN outburst in the nucleus B1.