

**Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
CBAT@CFA.HARVARD.EDU (science)
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304
Phone 617-495-7440/7244/7444 (for emergency use only)

V1663 AQUILAE = POSSIBLE NOVA IN AQUILA

M. Dennefeld and F. Riquebourg, Institut d'Astrophysique de Paris; and Y. Damerdjji, Observatoire de Haute-Provence (OHP), write that they obtained a calibrated spectrum (range 465–830 nm, resolution 0.6 nm) on June 11.1 UT of the possible nova reported on *IAUC* 8540, using the 1.93-m telescope (+ Carelec spectrograph) at OHP. The spectrum shows a very red continuum but no sign of molecular absorption bands, so reddening appears to be present; there are, however, various metallic stellar absorption features present, compatible with an early K-type star, so this may be a symbiotic nova (though the term 'peculiar nova' is perhaps more appropriate). There are also strong emission lines with narrow P-Cyg profiles at H α and O I (777 nm); the expansion velocities from these lines are 700 km/s in both cases (suggestive of a slow nova). Faint H β and numerous faint Fe II lines, all with P-Cyg profiles, are also seen in the range 480–640 nm. Due to the peculiar nature of this nova, continuous photometric and spectroscopic observations are encouraged.

N. N. Samus and E. Kazarovets, Institute of Astronomy, Russian Academy of Sciences, inform us that the designation V1663 Aql has been given to this nova (cf. *IAUC* 8540).

SUPERNOVA 2005cl IN MCG -01-53-20

Further to *IAUC* 8542, H. Pugh and W. Li report the LOSS discovery of an apparent supernova on unfiltered KAIT images taken on June 12.48 (at mag 18.3) and 14.45 UT (mag 18.2). An image taken on June 2.47 also showed a marginal detection of the new object (limiting mag \sim 18.8). SN 2005cl is located at $\alpha = 21^{\text{h}}02^{\text{m}}02^{\text{s}}.35$, $\delta = -6^{\circ}17'35''.7$ (equinox 2000.0), which is 18".0 west and 13".3 north of the nucleus of MCG -01-53-20. KAIT images taken on May 12.50 (limiting mag \sim 19.5) and 23.49 (limiting mag \sim 19.0) showed nothing at this position.

M. Modjaz, R. Kirshner, and P. Challis, Harvard-Smithsonian Center for Astrophysics, report that a spectrum (range 350–740 nm) of SN 2005cl, obtained on June 13.43 UT by P. Berlind with the F. L. Whipple Observatory 1.5-m telescope (+ FAST), shows it to be a type-II_n supernova. The spectrum consists of a blue continuum, indicating young age, and Balmer emission lines with a narrow H α component of \sim 500 km/s (FWHM) superposed on a weaker and broader component of \sim 4000 km/s (FWHM).