

**Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
CBAT@CFA.HARVARD.EDU (science)
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304
Phone 617-495-7440/7244/7444 (for emergency use only)

NOVA SCORPII 2005

M. L. Sitko and R. Kimes, University of Cincinnati; D. K. Lynch, R. W. Russell, and D. L. Kim, The Aerospace Corporation; and D. Griep, Infrared Telescope Facility (IRTF), report 3–14- μm spectroscopy of this object on July 27.375 UT using the instrument BASS on the IRTF. The spectrum shows a smooth, featureless continuum consistent with a Rayleigh-Jeans tail of a Planck function. The magnitude was ≈ 4.8 in the bands *L*, *M*, and *N*.

Visual magnitude estimates: July 26.735 UT, 8.9 (T. Cooper, Bredell, Kempton Park, S. Africa); 26.881, 8.9 (J. G. de S. Aguiar, Campinas, Brazil); 27.228, 9.0 (M. Linnolt, Woodside, CA).

SUPERNOVA 2005dc IN NGC 7107

P. Luckas, Perth, W. Australia; O. Trondal, Oslo, Norway; and M. Schwartz, Patagonia, AZ, report the discovery of an apparent supernova (mag 15.7) on an unfiltered CCD frames taken with a 0.35-m Tenagra telescope at Perth on July 26.70 and 27.62 UT in the course of the Tenagra Observatory Supernova Search. SN 2005dc is located at $\alpha = 21^{\text{h}}42^{\text{m}}22^{\text{s}}.66$, $\delta = -44^{\circ}47'21''.2$ (equinox 2000.0), which is $41''$ west and $9''.6$ north of NGC 7107 (the galaxy center having position end figures $26^{\text{s}}50$, $30''.8$). Nothing was visible at the position of 2005dc on an image taken by Luckas on July 5.61 (limiting magnitude ~ 18.5).

SUPERNOVA 2005da IN UGC 11301

M. Modjaz, P. Challis, and R. Kirshner, Harvard-Smithsonian Center for Astrophysics; and T. Matheson, National Optical Astronomy Observatory, report that a spectrogram (range 350–740 nm) of SN 2005da (cf. *IAUC* 8570), obtained by P. Berlind on July 26.26 UT with the F. L. Whipple Observatory 1.5-m telescope (+ FAST), reveals it to be a supernova, possibly a peculiar type-Ic supernova. No obvious lines of hydrogen or helium are detected. The spectrum consists of wide absorption features, similarly to peculiar type-Ic supernovae 1998bw and 2002ap (Foley *et al.* 2003, *PASP* **115**, 1220). In contrast to the spectra of supernovae 1998bw and 2002ap, this new object's spectrum also exhibits a wide absorption trough at 585 nm, having removed the NED recession velocity of 4495 km/s for the host galaxy (from Falco *et al.* 1999, *PASP* **111**, 438), which could be identified with Na I. There are peaks near 530 and 640 nm and absorptions near 470, 630, and 680 nm.