

**Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
CBAT@CFA.HARVARD.EDU (science)
URL <http://cfa-www.harvard.edu/iau/cbat.html> ISSN 0081-0304
Phone 617-495-7440/7244/7444 (for emergency use only)

NOVA OPHIUCHI 2006

G. Pojmanski, Warsaw University Astronomical Observatory, reports the discovery on ASAS-3 200-mm-f.l.-lens telephoto images (cf. *IAUC* 8617) of an apparent nova located at $\alpha = 17^{\text{h}}33^{\text{m}}13^{\text{s}}$, $\delta = -24^{\circ}21'1''$ (equinox 2000.0). Available ASAS *V* magnitudes: 2005 Oct. 26.013 UT, [14; 2006 Feb. 9.390, 12.08; 10.327, 12.04; 10.329, 11.94; 10.342, 12.07; 11.334, 11.48; 12.329, 11.07; 13.336, 11.47. The object is also clearly visible in the ASAS-3 *I*-band images taken on Feb. 8.379. H. Yamaoka, Kyushu University, provides his average of measurements of the position of the new nova from *I_c*-band CCD images taken by S. Kiyota on Feb. 12.83 and by H. Maehara on Feb. 10.83: $\alpha = 17^{\text{h}}33^{\text{m}}13^{\text{s}}.06$, $\delta = -24^{\circ}21'07''.1$ (rms $\sim 0''.2$; UCAC2 reference stars); the nearest star in the USNO-B1.0 catalog is $\sim 3''$ away from this position, and a red Digitized Sky Survey image from 1991 Aug. 11 does not show any object to limiting mag ~ 18 (affected by nearby stars). Yamaoka adds that low-resolution spectra of this new object — taken around Feb. 12.9 by K. Ayani with the Bisei Astronomical Observatory (BAO) 1.01-m telescope, by H. Naito and S. Narusawa (Nishi-Harima Astronomical Observatory) with the 2.0-m NAYUTA telescope, and by M. Fujii (Ibara, Okayama, Japan) with a 0.28-m telescope — show prominent Balmer lines with P-Cyg profiles superimposed on a very red continuum, which suggests that the object is a classical nova near maximum. Deep interstellar Na D absorption is also seen. From the BAO spectrum, the FWHM of H α emission is 560 km/s, and an absorption minimum of that is blueshifted by 530 km/s from the emission peak. O I 777.3-nm and Fe II emissions are also detected with P-Cyg profiles.

RS OPHIUCHI

This recurrent nova is in outburst for the first time since 1985 (cf. *IAUC* 4030, 4031), as indicated by the following visual magnitude estimates: Feb. 12.829 UT, 4.5 (H. Narumi, Ehime, Kita-gun, Japan; communicated by K. Hirosawa, Aichi, Inazawa-shi, Japan); 12.847, 4.6 (K. Kanai, Gunma, Isezaki-shi, Japan; via Hirosawa); 12.862, 4.4 (Kanai); 13.163, 4.8 (W. Renz, Karlsruhe, Germany); 13.27, 4.9 (A. Pereira, Cabo da Roca, Portugal). S. Nakano (Sumoto, Japan) forwards unfiltered CCD magnitudes of RS Oph from Y. Nakamura (Kameyama, Mie-ken, Japan; survey frames with a 105-mm *f*/2.8 lens): Feb. 9.855, 10.5; 12.8426, 4.5. ASAS *V* magnitudes from Pojmanski: 2005 Oct. 26.998, 10.72; 2006 Feb. 13.341, 5.4 (saturated); 13.344, 5.3.