

**Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
CBAT@CFA.HARVARD.EDU (science)
URL <http://www.cfa.harvard.edu/iau/cbat.html> ISSN 0081-0304
Phone 617-495-7440/7244/7444 (for emergency use only)

COMET C/2006 P1 (McNAUGHT)

C. M. Lisse and N. Dello Russo, Applied Physics Laboratory, Johns Hopkins University; Y. Fernandez, University of Central Florida; G. H. Jones, Mullard Space Science Laboratory, University College, London; and M. Sitko, Space Science Institute, report that the Spitzer Space Telescope's IRS instrument observed comet C/2006 P1 on May 4–5 (at $r = 2.4$ AU, $\Delta = 2.2$ AU). A very round, featureless coma with no obvious extension in the anti-sun direction appeared as the central source of emission in the peak-up imager at $16\ \mu\text{m}$. The 5- to $35\text{-}\mu\text{m}$ spectrum of the outflowing dust showed only a mild excess (~ 10 percent) due to silicate emission at $8\text{--}13\ \mu\text{m}$. The flux density at $10\ \mu\text{m}$ was ~ 0.1 Jy, and at $20\ \mu\text{m}$ was 0.6 Jy. The effective temperature of the dust was 190 ± 10 K. The local equilibrium temperature at 2.4 AU was 182 K. Lisse *et al.* estimate a production rate of dust to be $\sim 6 \times 10^3$ kg/s. The spectrum is remarkably featureless and dominated by infrared emission from large particles, and is similar to that derived from comet-surface mantles. This is unexpected for a comet that, in Dec. 2006–Jan. 2007, had demonstrated large outflows of material, a highly structured dust tail due to the presence of $0.1\text{--}10\text{-}\mu\text{m}$ dust particles, and was still emitting dust at the time of Spitzer observations at a rate comparable to the strongly-mid-infrared-featured comets C/1996 B2 (Hyakutake) and 1P/1982 U1 (Halley) when passing closest to the earth. Lisse *et al.* further surmise that either the comet has a very thick surface mantle that was only temporarily breached during the perihelion passage by a jet or the material being emitted in May 2007 is from a surface mantle that has regrown since the comet's perihelion passage four months earlier.

NOVA VULPECULAE 2007

C. Buil, Castanet, France, writes that a spectrogram of the possible nova reported on *IAUC* 8861, obtained on Aug. 9.91 UT with a 0.28-m telescope (+ Lhires spectrograph; resolution 6800 at $\text{H}\alpha$), shows evident $\text{H}\alpha$ with a deep P-Cyg profile and a FWHM of 1750 km/s (± 80 km/s); the intensity of the $\text{H}\alpha$ peak intensity is 2.4 times that of the local 670-nm continuum. M. Fujii, Kurashiki, Okayama, Japan, reports that a low-dispersion spectrogram (range $375\text{--}833$ nm; resolution ~ 1 nm) of the apparent nova was obtained on Aug. 9.48 UT with a 0.28-m reflector; the spectrum shows $\text{H}\alpha$, $\text{H}\beta$, $\text{H}\gamma$, and He I 447.2- , 471.3- , 492.2- , 501.5- , 587.6- , 667.8- , and 706.5-nm emissions with P-Cyg profiles ($\text{H}\alpha$ FWHM = 1900 km/s). These emissions suggest that the variable is indeed a classical nova.