

Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
 IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
 CBAT@CFA.HARVARD.EDU (science)
 URL <http://www.cfa.harvard.edu/iau/cbat.html> ISSN 0081-0304
 Phone 617-495-7440/7244/7444 (for emergency use only)

COMET 17P/HOLMES

C. Salyk and G. A. Blake, California Institute of Technology; M. J. Mumma, B. P. Bonev, M. A. DiSanti, and G. L. Villanueva, Goddard Space Flight Center; Y. Radeva, University of Maryland; K. Magee-Sauer, Rowan University; and E. L. Gibb, University of Missouri, St. Louis, report that they observed comet 17P/Holmes on Oct. 29 and 30 UT ($r = 2.46$ AU) using NIRSPEC at the W. M. Keck Observatory. On Oct. 29, H₂O (eleven lines, 2.9- μ m hot bands), C₂H₆ (eight Q-branches, 3.35- μ m ν_7 band), C₂H₂ (four lines, 3.0- μ m ν_3 band), HCN (nine lines, 3.0- μ m ν_3 band), and CH₃OH (3.52- μ m P-, Q-, R-branches, ν_3 band) were detected. Rotational temperatures for H₂O, HCN, and C₂H₆ were ~ 60 K. All lines were strongly peaked on the nucleus region, as expected for parent volatiles. Preliminary production rates ($\times 10^{27}$ molecules/s) are: H₂O, 275; C₂H₆, 5.9; C₂H₂, 1.5; HCN, 1.5; and CH₃OH, 11. The abundance ratios are: H₂O : C₂H₆ : C₂H₂ : HCN : CH₃OH = 100 : 2.2 : 0.54 : 0.54 : 4.0. Comet 17P seems to be enriched in organics, like C/2001 A2. On Oct. 30, abundance ratios for H₂O, C₂H₂, and HCN were similar to those on Oct. 29. The R0 and P2 lines of CO (4.7- μ m 1-0 band) were detected; if $T_{\text{rot}} = 60$ K, the formal CO abundance is 14 (± 4) relative to water (100).

COMETS C/2007 M5–M9 (SOHO)

Further to *IAUC* 8889, additional near-sun comets have been found on SOHO website images — C/2007 M5 and C/2007 M8 being non-group, and the rest being Kreutz sungrazers. C/2007 M5, which has the smallest perihelion distance on record ($q = 0.0011$ AU), entered the C2 field-of-view at peak brightness (mag ~ 6.5) and faded rapidly, disappearing from view before it reached the occulting disk. C/2007 M6 was stellar in appearance (mag ~ 7) in C3 images, but diffuse with no tail in C2 images. C/2007 M7 and C/2007 M9 were faint (mag ~ 8) and diffuse. C/2007 M8 was very small, circular (nearly stellar in appearance), and faint (mag ~ 7.5 –8), appearing to be marginally diffuse.

Comet	2007 UT	α_{2000}	δ_{2000}	Inst.	F	MPEC
C/2007 M5	June 25.229	6 ^h 06.0 ^m	+22°37'	C2	BZ	2007-U15
C/2007 M6	25.346	6 05.4	+21 38	C3/2	BZ	2007-U15
C/2007 M7	25.413	6 07.9	+21 39	C2	BZ	2007-U16
C/2007 M8	25.564	6 17.8	+22 32	C2	HS	2007-U16
C/2007 M9	27.604	6 16.5	+21 44	C2	BZ	2007-U16