

Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)
CBAT@CFA.HARVARD.EDU (science)
URL <http://www.cfa.harvard.edu/iau/cbat.html> ISSN 0081-0304
Phone 617-495-7440/7244/7444 (for emergency use only)

1998 WV₂₄

S. D. Benecchi, K. S. Noll, and E. A. Barker, Space Telescope Science Institute; W. M. Grundy, Lowell Observatory; and H. F. Levison, Southwest Research Institute, report that the transneptunian object 1998 WV₂₄ (cf. *MPECs* 1998-X12 and 1999-V06) is a binary. The observations were made during 2007 Aug. 27.1069–27.1375 UT with the Planetary Camera of the Wide Field Planetary Camera 2 on the Hubble Space Telescope, using the F606W filter (wide *V*) with one 260-s exposure at four dithered positions on the detector. The two components were separated by an angular distance of $0''.051 \pm 0''.002$, with the secondary fainter by 0.3 magnitude. The secondary was located at $0''.033 \pm 0''.003$ in α and $0''.039 \pm 0''.003$ in δ relative to the primary. Component centroids were determined by fitting the point-spread function.

V2468 CYGNI

D. K. Lynch, R. W. Russell, and R. J. Rudy, The Aerospace Corporation; M. Sitko, Space Science Institute; and C. E. Woodward, University of Minnesota, report 0.8- to 5.5- μm spectroscopy of the nova V2468 Cyg (cf. *IAUC* 8927, 8928, 8936) on Oct. 4 UT using the Infrared Telescope Facility (+ Spex). Though V2468 Cyg had faded by more than a factor of two since mid-August, the coronal lines have strengthened significantly. The [Si VI] line at 1.96 μm is among the strongest of the emission features. Lines of [S VIII], [S IX], [Al IX], and [Ca VIII] are also present. Several Rydberg lines are present, including those at 0.89, 1.19, 1.11, 1.55, and 2.1 μm .

V2670 OPHIUCHI

M. Sitko, Space Science Institute; D. K. Lynch, R. W. Russell, and R. J. Rudy, The Aerospace Corporation; and C. E. Woodward, University of Minnesota, report 0.8- to 5.5- μm spectroscopy of the nova V2670 Oph (cf. *IAUC* 88947, 8948, 8949, 8956) on Oct. 4 UT using the Infrared Telescope Facility (+ Spex). The spectrum remains low in excitation, still displaying emission lines of Fe II, N I, and O I. However, the C I lines that were strong in mid-June are barely detectable, and the He I 1.083- μm line is now the strongest feature in the infrared spectrum. The broad components of the emission lines have disappeared entirely. This nova has not formed dust and is unlikely to do so at this late stage.